

MENIIT

NEET | IIT-JEE | FOUNDATION

Corporate Office: 44-A/1, Kalu Sarai, New Delhi 110016 | Web: www.meniit.com

JEE Advanced : Paper-2 (2011)

IMPORTANT INSTRUCTIONS

A. General:

1. The **question paper CODE** is printed on the right hand top corner of this sheet and on the back page (page no. 18) of this booklet.
2. No additional sheets will be provided for rough work.
3. Blank papers, clipboards, log tables, slide rules, calculators, cellular phones, pagers, and electronic gadgets in any form are not allowed.
4. Write your name and registration number in the space provided on the back page of this booklet.
5. The answer sheet, a machine-gradable. Objective Response sheet (**ORS**), is provided separately
6. **DO NOT TAMPER WITH /MUTILATE THE ORS OR THE BOOKLET.**
7. Do not break the seals of the question paper booklet before instructed to do so by the invigilators.
8. This question paper contains 18 pages having 69 questions.
9. On breaking the seals please check that all the questions are legible.

B. Filling the Right Part of the ORS:

10. The ORS also has a **CODES** printed on its lower and upper parts.
11. Make sure the **CODE** on the **ORS** is the same as that on this booklet. If the Codes do not match, ask **for a change of the Booklet.**
12. Write your Registration No., Name and Name of centre and sign with pen in appropriate boxes. Do not write these anywhere else. Darken the appropriate bubbles under each digit of your Registration No. with HB Pencil.

C. Question paper format and Marking scheme:

13. The question paper consists of **3 parts** (Chemistry, Physics and Mathematics). Each part consists of four sections.
14. In **Section I** (Total marks : 24), for each question you will be awarded **3 marks** if you darken **ONLY** the bubble corresponding to the correct answer and **zero mark** if no bubbles are darkened. In all other cases, **minus one (-1) mark** will be awarded.
15. In **Section II** (Total Marks : 16), for each question you will be awarded **4 marks** if you darken all the bubble(s) corresponding to the correct answer(s) **ONLY** and **zero mark** otherwise. There are **no negative marks** in this section.
16. In **Section III** (Total Marks : 24), for each question you will be **awarded 4 marks** if you darken **only** the bubble corresponding to the correct answer and **zero marks otherwise. There are no negative marks** in this section.
17. In **Section IV** (Total Marks : 16), for each question you will be **awarded 2 marks** for each row in which you have darkened all the bubble(s) corresponding to the correct answer(s) **ONLY** and **zero marks** otherwise. Thus each question in this section carries a **maximum of 8 marks**. There are **no negative** marks in this section.





PART A : CHEMISTRY

SECTION -I (Total Marks : 24)


(Single Correct Answer Type)

This section contains 8 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONLY ONE is correct.


1. Amongst the compounds given, the one that would form a brilliant colored dye on treatment with NaNO_2 in dil. HCl followed by addition to an alkaline solution of β -naphthol is :



Ans. C



2. The major product of the following reaction is :




(A) a hemiacetal

(B) an acetal


(C) an ether

(D) an ester

Ans. B



3. The following carbohydrate is



(A) a ketohexose

(B) an aldohexose

(C) an α -furanose

(D) an α -pyranose

Ans. B

Sol. Aldohexose

4. Oxidation states of the metal in the minerals haematite and magnetite, respectively, are
 (A) II, III in haematite and III in magnetite (B) II, III in haematite and II in magnetite
 (C) II in haematite and II, III in magnetite (D) III in haematite and II, III in magnetite

Ans. D

Sol. Haematite is $\overset{+3}{\text{Fe}}_2\text{O}_3$

Magnetite is Fe_3O_4 or $\overset{+2}{\text{Fe}}\text{O}.\overset{+3}{\text{Fe}}_2\text{O}_3$

5. Among the following complexes (**K–P**)

$\text{K}_3[\text{Fe}(\text{CN})_6]$ (**K**), $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$ (**L**), $\text{Na}_3[\text{Co}(\text{oxalate})_3]$ (**M**), $[\text{Ni}(\text{H}_2\text{O})_6]\text{Cl}_2$ (**N**), $\text{K}_2[\text{Pt}(\text{CN})_4]$ (**O**) and $[\text{Zn}(\text{H}_2\text{O})_6](\text{NO}_3)_2$ (**P**)

the diamagnetic complexes are :

- (A) K, L, M, N (B) K, M, O, P (C) L, M, O, P (D) L, M, N, O

Ans. C

Sol. (L) : $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$

(M) : $\text{Na}_3[\text{Co}(\text{Ox})_3]$

(O) : $\text{K}_2[\text{Pt}(\text{CN})_4]$

(P) : $[\text{Zn}(\text{H}_2\text{O})_6](\text{NO}_3)_2$


6. Passing H_2S gas into a mixture of Mn^{2+} , Ni^{2+} , Cu^{2+} and Hg^{2+} ions in an acidified aqueous solution precipitates

- (A) CuS and HgS (B) MnS and CuS (C) MnS and NiS (D) NiS and HgS

Ans. A

Sol. Cu^{+2} , Hg^{+2} are group II basic radicals

7. Consider the following cell reaction:



At $[\text{Fe}^{2+}] = 10^{-3} \text{ M}$, $P(\text{O}_2) = 0.1 \text{ atm}$ and $\text{pH} = 3$, the cell potential at 25°C is :

- (A) 1.47 V (B) 1.77 V (C) 1.87 V (D) 1.57 V

Ans. D

Sol. $E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{.0591}{4} \log \frac{[\text{Fe}^{+2}]}{[\text{Po}_2][\text{H}^+]^4}$

$$= 1.64 - \frac{.591}{4} \log \frac{[10^{-3}]^2}{[.1][10^{-3}]}$$

$$= 1.57 \text{ V}$$

8. The freezing point (in °C) of a solution containing 0.1 g of $K_3[Fe(CN)_6]$ (Mol. Wt. 329) in 100 g of water ($K_f = 1.86 \text{ K kg mol}^{-1}$) is
- (A) -2.3×10^{-2} (B) -5.7×10^{-2} (C) -5.7×10^{-3} (D) -1.2×10^{-2}

Ans. A

Sol. $\Delta T = k_f \times m \times i \times 1000$

$$= 1.86 \times \frac{0.1}{329 \times 100} \times 4 \times 1000$$

$$= 2.26 \times 10^{-2} \approx 2.3 \times 10^{-2}$$

SECTION-II (Total Marks : 16)

(Multiple Correct Answer(s) Type)

This section contains 4 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONE OR MORE may be correct.

9. The equilibrium



in aqueous medium at 25°C shifts towards the left in the presence of :

- (A) NO_3^- (B) Cl^- (C) SCN^- (D) CN^-

Ans. B, C, D


Sol. $\text{Cu}_2 \text{Cl}_2$, $\text{Cu}_2(\text{CN})_2$ and $\text{Cu}_2(\text{SCN})_2$ are stable

10. Reduction of the metal centre in aqueous permanganate ion involves :


- (A) 3 electrons in neutral medium (B) 5 electrons in neutral medium
(C) 3 electrons in alkaline medium (D) 5 electrons in acidic medium

Ans. A, C, D

Sol. \rightarrow In alkaline solution, KMnO_4 is first reduced to manganate and then to insoluble MnO_2



11. The correct functional group X and the reagent/reaction conditions Y in the following scheme are :




- (A) $\text{X} = \text{COOCH}_3$, $\text{Y} = \text{H}_2 / \text{Ni} / \text{heat}$ (B) $\text{X} = \text{CONH}_2$, $\text{Y} = \text{H}_2 / \text{Ni} / \text{heat}$
(C) $\text{X} = \text{CONH}_2$, $\text{Y} = \text{Br}_2 / \text{NaOH}$ (D) $\text{X} = \text{CN}$, $\text{Y} = \text{H}_2 / \text{Ni} / \text{heat}$

Ans. A, B, C, D

Sol. Factual

* The most appropriate answer to this question is (A,B,C,D)
But because of ambiguity in language, IIT has declared (C & D)
as correct answer

12. For the first order reaction



- (A) the concentration of the reactant decreases exponentially with time.
(B) the half-life of the reaction decreases with increasing temperature
(C) the half-life of the reaction depends on the initial concentration of the reactant
(D) the reaction proceeds to 99.6 % completion in eight half-life duration.

Ans. A, B, D

Sol. $C_A = C_{A_0} e^{-kt}$ [A]

$$t_{\frac{1}{2}} = \frac{0.693}{K} = \frac{0.693}{A_0 e^{-E_a/RT}} = \frac{0.693}{A_0} e^{E_a/RT}$$
 [B]


$$\frac{0.4}{100} = \left(\frac{1}{2}\right)^4 = \frac{4}{100} = n = \frac{\log\left(\frac{4}{100}\right)}{\log\left(\frac{1}{2}\right)} = 8$$
 [D]

SECTION-III (Total Marks : 24)

(Integer Answer Type)

This Section contains a group of 6 questions. The answer to each questions is a single digit integer ranging from 0 to 9. The bubble corresponding to the correct answer is to be darkened in the ORS.


13. The total number of contributing structures showing hyperconjugation (involving C–H bonds) for the following carbocation is :



Ans. 6



Sol. 6 ($\alpha - \text{H} \rightarrow 6$)

14. Among the following the number of compounds than can react with PCl_5 to give POCl_3 is :



Ans. 4

Sol. $\text{PCl}_5 + \text{H}_2\text{O} \longrightarrow \text{POCl}_3 + 2\text{HCl}$



15. The volume (in mL) of 0.1 M AgNO_3 required for complete precipitation of chloride ions present in 30 mL of 0.01 M solution of $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2$, as silver chloride is close to :

Ans. 6

Sol. $0.1V = 30 \times 0.01 \times 2$

$$v = \frac{0.3 \times 2}{0.1} = 6 \text{ ml}$$

16. In 1 L saturated solution of AgCl [$K_{\text{sp}}(\text{AgCl}) = 1.6 \times 10^{-10}$], 0.1 mol of CuCl [$K_{\text{sp}}(\text{CuCl}) = 1.0 \times 10^{-6}$] is added. The resultant concentration of Ag^+ in the solution is 1.6×10^{-x} . The value of "x" is :

Ans. 7

Sol. $[\text{Ag}^+] = \frac{K_1}{\sqrt{K_1 + K_2}} \because K_1 \ll K_2 \therefore K_1 + K_2 \cong K_2$


$$\therefore [\text{Ag}^+] = \frac{1.6 \times 10^{-10}}{\sqrt{1.0 \times 10^{-6}}} = 1.6 \times 10^{-7}$$

$$x = 7$$

17. The number of hexagonal faces that are present in a truncated octahedron is


Ans. 8

Sol.




8 Hexagonal faces

18. The maximum number of isomers (including stereoisomers) that are possible on mono-chlorination of the following compound, is :



Ans. 8



SECTION-IV (Total Mark : 16)

(Matric-Match Type)

This section contains 2 questions. Each question has four statements (A,B,C and D) given in Column I and five statements (p, q, r, s and t) in Column II. Any given statement in Column I can have correct matching with ONE OR MORE statement(s) in Column II. For example, if for a given question, statement B matches with the statements given in q and r, then for that particular question, against statement B, darken the bubbles corresponding to q and r in the ORS.

19. Match the reactions in column I with appropriate types of steps / reactive intermediate involved in these reactions as given in column II

Column I		Column II
(A)	$\xrightarrow{\text{aq. NaOH}}$	(p) Nucleophilic substitution
(B)	$\xrightarrow{\text{CH}_3\text{MgI}}$	(q) Electrophilic substitution
(C)	$\xrightarrow{\text{H}_2\text{SO}_4}$	(r) Dehydration
(D)	$\xrightarrow{\text{H}_2\text{SO}_4}$	(s) Nucleophilic addition
		(t) Carbanion

Ans. (A) r,s,t (B) p,s,t (C) r,s (D) q,r

Sol. Factual

* The most appropriate answer to this question is





A → r, s, t; B → p, s, t; C → r, s; D → q, r

But because of ambiguity in language, IIT has declared

A → r, s, t; B → p, s; C → r, s; D → q & r as correct answer

20. Match the transformations in column I with appropriate options in column II

Column I



Column II

(p) phase transition

(q) allotropic change

(r) ΔH is positive


(s) ΔS is positive

(t) ΔS is negative


Ans. (A) p,r,s (B) r,s (C) t (D) q,t

Sol. [A] $\text{CO}_2(\text{s}) \longrightarrow \text{CO}_2(\text{g})$

p, r, s



r, s



t



p, q, t


PART B : PHYSICS

SECTION-I

Single Correct Choice Type

This section contains 8 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONLY ONE is correct.


21. A ball of mass 0.2 kg rests on a vertical post of height 5 m. A bullet of mass 0.01 kg, travelling with a velocity V m/s in a horizontal direction, hits the centre of the ball. After the collision, the ball and bullet travel independently. The ball hits the ground at a distance of 20 m and the bullet at a distance of 100 m from the foot of the post. The initial velocity V of the bullet is :



- (A) 250 m/s (B) $250\sqrt{2}$ m/s (C) 400 m/s (D) 500 m/s

Ans. D

Sol. $m_1 = 0.01$ kg $m_2 = 0.2$ kg



$$T = \sqrt{\frac{2H}{g}} = 1 \text{ sec}$$

Let v_1 & v_2 be velocity of bullet & ball respectively just after collision.

$$v_2 \times 1 = 20 \Rightarrow v_2 = 20$$

$$\& v_1 = 100$$

From conservation of momentum

$$0.01 \times v = (0.01 \times 100) + (0.2 \times 20)$$

$$0.01 v = 1 + 4 = 5$$

$$v = \frac{5}{10^{-2}} = 500 \text{ m/sec.}$$

22. The density of a solid ball is to be determined in an experiment. The diameter of the ball is measured with a screw gauge, whose pitch is 0.5 mm and there are 50 divisions on the circular scale. The reading on the main scale is 2.5 mm and that on the circular scale is 20 divisions. If the measured mass of the ball has a relative error of 2%, the relative percentage error in the density is :
- (A) 0.9% (B) 2.4% (C) 3.1% (D) 4.2%

Ans. C

Sol. Pitch = 0.5 mm
divisions on the = 50
circular scale

$$\Rightarrow \text{least count of screw gauge} = \frac{0.5}{50} = 0.01$$

main scale, reading = 2.5 mm

circular scale reading = 20


$$\Rightarrow \text{reading} = 2.5 \text{ mm} + (20 \times 0.01) \text{ mm} \\ = 2.5 \text{ mm} + 0.2 \text{ mm} = 2.7 \text{ mm}$$

$$\rho = \frac{m}{\frac{4\pi}{3} \left[\frac{D}{2} \right]^3}$$

$$\frac{\Delta\rho}{\rho} = \frac{\Delta m}{m} + 3 \frac{\Delta D}{D}$$

$$\% \text{ error} = \frac{\Delta\rho}{\rho} \times 100 = 2\% + 3 \left(\frac{0.01}{2.7} \right) \times 100 = 3.1.$$

23. A wooden block performs SHM on a frictionless surface with frequency, ν_0 . The block carries a charge +Q on its surface. If now a uniform electric field \vec{E} is switched on as shown, then the SHM of the block will be :




- (A) of the same frequency and with shifted mean position.
(B) of the same frequency and with the same mean position.
(C) of changed frequency and with shifted mean position.
(D) of changed frequency and with the same mean position.


Ans. A

Sol. In order to have net force zero, the mean position will be shifted towards right but the time period will remain unaffected.

24. A light ray traveling in glass medium is incident on glass-air interface at an angle of incidence θ . The reflected (R) and transmitted (T) intensities, both as function of θ , are plotted. The correct sketch is :




Ans. C



When $\theta > \theta_c$, no ray will transmit

$\Rightarrow T = 0, T + R = 100\%$ and $R > 0$




25. A satellite is moving with a constant speed 'V' in a circular orbit about the earth. An object of mass 'm' is ejected from the satellite such that it just escapes from the gravitational pull of the earth. At the time of its ejection, the kinetic energy of the objects :

- (A) $\frac{1}{2}mV^2$ (B) mV^2 (C) $\frac{3}{2}mV^2$ (D) $2mV^2$

Ans. B

Sol.




$$\Rightarrow \frac{mv^2}{r} = \frac{GmM_e}{r^2} \Rightarrow r = \frac{GM_e}{v^2} \dots(1)$$

If K.E. of mass m = was k then from

$$E = K - \frac{GmM_e}{r} = 0 \Rightarrow K = m \left(\frac{GM_e}{r} \right) = mv^2$$

26. A long insulated copper wire is closely wound as a spiral of 'N' turns. The spiral has inner radius 'a' and outer radius 'b'. The spiral lies in the X-Y plane and a steady current 'I' flows through the wire. The Z-component of the magnetic field at the centre of the spiral is :



- (A) $\frac{\mu_0 N I}{2(b-a)} \ln \left(\frac{b}{a} \right)$ (B) $\frac{\mu_0 N I}{2(b-a)} \ln \left(\frac{b+a}{b-a} \right)$ (C) $\frac{\mu_0 N I}{2b} \ln \left(\frac{b}{a} \right)$ (D) $\frac{\mu_0 N I}{2b} \ln \left(\frac{b+a}{b-a} \right)$

Ans. A


Sol. $(b - a) \rightarrow N$

$$dr \rightarrow \left(\frac{N}{b-a} \right) dr$$

$$B = \frac{\mu_0 I N}{2(b-a)} \ln(b/a)$$

$$dB = \left(\frac{\mu_0 I}{2r} \right) \left(\frac{N}{b-a} \right) dr$$

$$B = \frac{\mu_0 I N}{2(b-a)} \int_a^b \frac{dr}{r}$$




27. A point mass is subjected to two simultaneously sinusoidal displacements in x-direction $x_1(t) = A \sin \omega t$ and $x_2(t) = A \sin \left(\omega t + \frac{2\pi}{3} \right)$. Adding a third sinusoidal displacement $x_3(t) = B \sin (\omega t + \phi)$ brings the mass to a complete rest. The values of B and ϕ are :

- (A) $\sqrt{2}A, \frac{3\pi}{4}$ (B) $A, \frac{4\pi}{3}$ (C) $\sqrt{3}A, \frac{5\pi}{6}$ (D) $A, \frac{\pi}{3}$

Ans. B

Sol.




Here $\phi = \pi + \theta$

$$A \cos 30^\circ = B \sin \theta \Rightarrow B \sin \theta = \frac{\sqrt{3}A}{2} \text{ and } A \sin 30^\circ + B \cos \theta = A \Rightarrow B \cos \theta = \frac{A}{2}$$

Solving above, $B = A$ and $\theta = 60^\circ = \frac{\pi}{3}$. Hence $f = 240^\circ = \frac{4\pi}{3}$

28. Which of the field patterns given below is valid for electric field as well as for magnetic field.



Ans. C

Sol. Induced electric field lines.

Magnetic field lines due to wire carrying current.

SECTION-II

Multiple Correct Choice Type


This section contains 4 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONE OR MORE may be correct.

29. A series R-C circuit is connected to AC voltage source. Consider two cases ; (A) when C is without a dielectric medium and (B) when C is filled with dielectric of constant 4. The current I_R through the resistor and voltage V_C across the capacitor are compared in the two cases. Which of the following is/are true ?


- (A) $I_R^A > I_R^B$ (B) $I_R^A < I_R^B$ (C) $V_C^A > V_C^B$ (D) $V_C^A < V_C^B$

Ans. B,C

Sol.



$$Z_1 = \sqrt{R^2 + \left(\frac{1}{\omega C}\right)^2}$$




$$Z_2 = \sqrt{R^2 + \left(\frac{1}{\omega C}\right)^2}$$

$$Z_1 > Z_2 \quad \therefore \quad I_R^A < I_R^B$$

$$V_C^A = \frac{I_C^A}{\omega C} \quad ; \quad V_C^B = \frac{I_C^B}{\omega C}; \quad V_C^B < V_C^A$$

30. Two solid spheres A and B of equal volumes but of different densities d_A and d_B are connected by a string. They are fully immersed in a fluid of density d_f . They get arranged into an equilibrium state as show in the figure with a tension in the string. The arrangement is possible only if :



(A) $d_A < d_f$


(B) $d_B > d_f$

(C) $d_A > d_f$

(D) $d_A + d_B = 2d_f$

Ans. A,B,D


Sol.



system will be in equilibrium with tension in string only if $d_f > d_A$ and $d_B > d_f$. If both A & B are considered as a system then

$$2Vd_f g = V(d_A + d_B)g \Rightarrow d_A + d_B = 2d_f$$

31. A thin ring of mass 2 kg and radius 0.5 m is rolling without slipping on a horizontal plane with velocity 1 m/s. A small ball of mass 0.1 kg, moving with velocity 20 m/s in the opposite direction, hits the ring at a height of 0.75 m and goes vertically up with velocity 10 m/s. Immediately after the collision :



- (A) the ring has pure rotation about its stationary CM
- (B) the ring comes to a complete stop.
- (C) friction between the ring and the ground is to the left.
- (D) there is no friction between the ring and the ground.

Ans. A,C

Sol. As no data is given about nature of horizontal plane.

* The most appropriate answer to this question is (A,C), but because of ambiguity in language, IIT has declared [(A, C), (A)] as correct answer

32. Which of the following statement(s) is/are correct?

- (A) if the electric field due to a point charge varies as $r^{-2.5}$ instead of r^{-2} , then the Gauss law will still be valid.
- (B) The Gauss law can be used to calculate the field distribution around an electric dipole.
- (C) If the electric field between two point charges is zero somewhere, then the sign of the two charges is the same.
- (D) The work done by the external force in moving a unit positive charge from point A at potential V_A to point B at potential V_B is $(V_B - V_A)$.

Ans. C,D

SECTION - III

Integer Answer Type

This section contains 6 questions. The answer to each of the questions is a single-digit integer, ranging from 0 to 9. The bubble corresponding to the correct answer is to be darkened in the ORS.

33. A silver sphere of radius 1 cm and work function 4.7 eV is suspended from an insulating thread in free-space. It is under continuous illumination of 200 nm wavelength light. As photoelectrons are emitted, the sphere gets charged and acquires a potential. The maximum number of photoelectrons emitted from the sphere is $A \times 10^Z$ (where $1 < A < 10$). The value of 'Z' is :

Ans. 7

Sol. Energy of photon $\approx \frac{12420}{200} \text{ eV} = 6.2 \text{ eV}$

Maximum KE of a electron = 6.2 eV – 4.7 eV

When potential on surface of sphere becomes equal to 1.5V

$$\frac{q}{4\pi\epsilon_0 r} = 1.5 \text{ V} \Rightarrow q = 1.5 \times (4\pi\epsilon_0) \times r$$

$$\text{No. of photoelectron emitted } n = \frac{1.5 \times (4\pi\epsilon_0) r}{1.6 \times 10^{-19}} = 1.04 \times 10^7$$

34. A train is moving along a straight line with a constant acceleration 'a'. A boy standing in the train throws a ball forward with a speed of 10 m/s, at an angle of 60° to the horizontal. The boy has to move forward by 1.15 m inside the train to catch the ball back at the initial height. The acceleration of the train, in m/s² is :

Ans. 5

Sol. $T = \frac{2 \times 10 \times \sqrt{3}}{2 \times 10} = \sqrt{3} \text{ sec}$


$x = 10 \cos 60^\circ (T) = 5\sqrt{3} \text{ m}$

In frame of train,

$5\sqrt{3} = \frac{1}{2} \times a \times (\sqrt{3})^2 + 1.15$ (a : acceleration of train)

$a = 5 \text{ m/sec}^2$

35. A block of mass 0.18 kg is attached to a spring of force-constant 2N/m. The coefficient of friction between the block and the floor is 0.1. Initially the block is at rest and the spring is un-stretched. An impulse is given to the block as shown in the figure. The block slides a distance of 0.06 m and comes to rest for the first time. The initial velocity of the block in m/s is $V = N/10$. Then N is :



Ans. 4


Sol. Using W – E theorem

$\frac{1}{2} \times m(u)^2 = \frac{1}{2} K(x)^2 + \mu mg(x)$


$\frac{1}{2} \times (0.18) u^2 = \frac{1}{2} \times 2 \times 36 \times 10^{-4} + 0.1 \times 0.18 \times 10 \times 0.06$

$\Rightarrow u = 0.4 \text{ m/sec.}$

$\Rightarrow \frac{4}{10} \text{ m/ sec.}$




36. Two batteries of different emfs and different internal resistances are connected as shown. The voltage across AB in volts is :



Ans. 5

Sol. $V_A - V_B = \frac{\frac{6}{1} + \frac{3}{2}}{\frac{1}{1} + \frac{1}{2}} = \frac{6+1.5}{1.5} = \frac{7.5}{1.5} = 5V$

- 37.** Water (with refractive index $= \frac{4}{3}$) in a tank is 18 cm deep. Oil of refractive index $\frac{7}{4}$ lies on water making a convex surface of radius of curvature 'R = 6 cm' as shown. Consider oil to act as a thin lens. An object 'S' is placed 24 cm above water surface. The location of its image is at 'x' cm above the bottom of the tank. Then 'x' is :



Ans. 2

Sol. $\frac{\mu_3}{v} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R_1} + \frac{\mu_3 - \mu_2}{R_2}$

$$\frac{4}{3V} - \frac{1}{-24} = \frac{\frac{7}{4} - 1}{6} + \frac{\frac{4}{3} - \frac{7}{4}}{\infty} \Rightarrow \frac{4}{3V} + \frac{1}{24} = \frac{1}{8} \Rightarrow \frac{4}{3V} = \frac{1}{12} \Rightarrow V = 16 \text{ cm}$$

∴ Ans. = (18 - 16) cm = 2 cm

- 38.** A series R-C combination is connected to an AC voltage of angular frequency $\omega = 500$ radian/s. If the impedance of the R-C circuit is $R\sqrt{1.25}$, the time constant (in millisecond) of the circuit is :

Ans. 4

Sol. $z = R\sqrt{1.25}$

$$\tau = RC$$

$$R^2 + \left(\frac{1}{500C}\right)^2 = Z^2$$

$$R^2 + \left(\frac{1}{500C}\right)^2 = R^2 \times 1.25$$

$$\left(\frac{1}{500C}\right)^2 = 0.25 R^2 \Rightarrow \frac{1}{500C} = 0.5 R$$

$$\frac{1}{2500} = RC$$

$$\frac{1}{250} = RC$$

0.004 sec = RC





RC = 4 mill second.

SECTION-IV

(Matrix-Match Type)

This section contains 2 questions. Each question has four statements (A, B, C and D) given in Column-I and five statements (p, q, r, s and t) in Column-II. Any given statement in Column-I can have correct matching with ONE or MORE statement(s) given in Column-II. For example, if for a given question, statement B matches with the statements given in q and r, then for the particular question, against statement B, darken the bubbles corresponding to q and r in the ORS.

39. Column-I shows four systems, each of the same length L, for producing standing waves. The lowest possible natural frequency of a system is called is fundamental frequency, whose wavelength is denoted as λ_f . Match each system with statements given in Column-II describing the nature and wavelength of the standing waves.

Column-I	Column-II
(A) Pipe closed at one end 	(p) Longitudinal waves
(B) Pipe open at both ends 	(q) Transverse waves
(C) Stretched wire clamped at both ends 	(r) $\lambda_f = L$
(D) Stretched wire clamped at both ends and at mid-point. 	(s) $\lambda_f = 2L$
	(t) $\lambda_f = 4L$

Ans. (A) p,t (B) p,s (C) q,s (D) q,r

Sol.




$$\frac{\lambda_f}{4} = L$$

$$\Rightarrow \lambda_f = 4L$$

(C) Stretched wire clamped at both ends

(B) Longitudinal waves



$$\frac{\lambda_f}{2} = L$$






$$\frac{\lambda_f}{2} + \frac{\lambda_f}{2} = L$$

$$\frac{\lambda_f}{2} = L \Rightarrow \lambda_f = 2L$$

$$\Rightarrow \lambda_f = L$$

40. One mole of a monatomic ideal gas is taken through a cycle ABCDA as shown in the P-V diagram. Column-II gives the characteristics involved in the cycle. Match them with each of the processes given in Column-I.



Column-I

- (A) Process A → B
- (B) Process B → C
- (C) Process C → D
- (D) Process D → A

Column-II

- (p) Internal energy decreases
- (q) Internal energy increases
- (r) Heat is lost
- (s) Heat is gained
- (t) Work is done on the gas

Ans. (A) p,r,t (B) p,r (C) q,s (D) r,t

Sol. Process AB : (Pressure is constant)

$$\text{If } T_A = T \Rightarrow T_B = T/3$$

$$\text{So } \Delta U = \text{Negative} [\because \Delta U = nC_V\Delta T]$$

$$\Delta W = nR\Delta T = \text{Negative}$$

$$\Delta Q = \Delta U + \Delta W = \text{Negative}$$

Process BC : (Volume is constant)

$$\text{If } T_B = \frac{T}{3} \text{ then } T_C = \frac{T}{9}$$

$$\Delta U = nC_V\Delta T = \text{Negative}$$

$$\Delta W = \text{Zero}$$

$$\Delta Q = \text{Negative}$$

Process C → D : (Pressure is constant)

$$\text{If } T_C = T/9 \text{ then } T_D = T$$

$$\Delta U = nC_V\Delta T = \text{positive}$$

$$\Delta W = \text{positive}$$

$\Delta Q = \text{positive}$

Process D \rightarrow A :

$T_D = T$ and $T_A = T$

Hence process is isothermal

$\Delta U = 0$

$\Delta W = \text{negative}$

$\Delta Q = \text{negative}$

MENIIT
NEET | IIT-JEE | FOUNDATION

PART C: MATHEMATICS

Section - I (Total Marks : 24)

(Single Correct Answer Type)

This section contains **8 multiple choice questions**. Each question has four choices (A), (B), (C) and (D) out of which **ONLY ONE** is correct.

41. Let $f : [-1, 2] \rightarrow [0, \infty]$ be a continuous function such that $f(x) = f(1-x)$ for all $x \in [-1, 2]$.

Let $R_1 = \int_{-1}^2 x f(x) dx$, and R_2 be the area of the region bounded by $y = f(x)$, $x = -1$, $x = 2$ and the x -axis.

Then :

- (A) $R_1 = 2R_2$ (B) $R_1 = 3R_2$ (C) $2R_1 = R_2$ (D) $3R_1 = R_2$

Ans. **C**

Sol. $R_1 = \int_{-1}^2 x f(x) dx$ (i)

$$R_1 = \int_{-1}^2 (1-x)(1-x) dx$$

$$= \int_{-1}^2 (1-x)f(x) dx$$
(ii)

(i) + (ii)

$$2R_1 = \int_{-1}^2 f(x) dx = R_2$$

$\therefore 2R_1 = R_2$

42. Let $f(x) = x^2$ and $g(x) = \sin x$ for all $x \in \mathbb{R}$. Then the set of all x satisfying

$(f \circ g \circ g \circ f)(x) = (g \circ g \circ f)(x)$, where $(f \circ g)(x) = f(g(x))$, is

- (A) $\pm\sqrt{n\pi}$, $n \in \{0, 1, 2, \dots\}$ (B) $\pm\sqrt{n\pi}$, $n \in \{1, 2, \dots\}$
 (C) $\frac{\pi}{2} + 2n\pi$, $n \in \{\dots, -2, -1, 0, 1, 2, \dots\}$ (D) $2n\pi$, $n \in \{\dots, -2, -1, 0, 1, 2, \dots\}$

Ans. **A**

Sol. $g \circ f(x) = g(f(x)) = g(x^2) = \sin x^2$
 $g \circ (g \circ f)(x) = g(\sin x^2) = \sin(\sin x^2)$
 $f \circ (g \circ g \circ f)(x) = f(\sin(\sin x^2)) = (\sin(\sin x^2))^2$
 $\therefore (\sin(\sin x^2))^2 = \sin(\sin x^2)$
 $\sin(\sin x^2) (\sin(\sin x^2) - 1) = 0$
 $\sin(\sin x^2) = 0$ or $\sin(\sin x^2) = 1$
 $\sin x^2 = n\pi$ $\sin x^2 = 2n\pi + \frac{\pi}{2}$


At $n = 0$	At $n = 0$
$\sin x^2 = 0$	$\sin x^2 = \frac{\pi}{2}$
$x^2 = n\pi$	Not possible
$x = \pm\sqrt{n\pi}$;	$n \in \{0, 1, 2, \dots\}$

43. Let (x, y) be any point on the parabola $y^2 = 4x$. Let P be the point that divides the line segment from $(0, 0)$ to (x, y) in the ratio 1 : 3. Then the locus of P is :

- (A) $x^2 = y$ (B) $y^2 = 2x$ (C) $y^2 = x$ (D) $x^2 = 2y$

Ans. C

Sol. $h = \frac{t^2}{4}, k = \frac{2t}{4}$
 $t^2 = 4h, y = 2k$
 so $4k^2 = 4h$
 $\therefore k^2 = h$



Hence required locus is $y^2 = x$

44. Let P (6, 3) be a point on the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$. If the normal at the point P intersects the x-axis at (9, 0), then the eccentricity of the hyperbola is :

- (A) $\sqrt{\frac{5}{2}}$ (B) $\sqrt{\frac{3}{2}}$ (C) $\sqrt{2}$ (D) $\sqrt{3}$

Ans. B

Sol. Equation of the normal at (6,3) is

$$\frac{a^2x}{6} + \frac{b^2y}{3} = a^2 + b^2$$

It passes through (9, 0)

$$\text{so } \frac{9a^2}{6} = a^2 + b^2$$

$$\Rightarrow b^2 = \frac{a^2}{2}$$

$$\therefore e^2 - 1 = \frac{1}{2}$$

$$e^2 = \frac{3}{2} \quad \Rightarrow e = \sqrt{\frac{3}{2}}$$

45. A value of b for which the equations

$$x^2 + bx - 1 = 0$$

$$x^2 + x + b = 0$$

have one root in common is

- (A) $-\sqrt{2}$ (B) $-i\sqrt{3}$ (C) $i\sqrt{5}$ (D) $\sqrt{2}$

Ans. B

Sol. $x^2 + bx - 1 = 0$ (i)

$x^2 + x + b = 0$ (ii)

(i) - (ii) we get $x = \frac{b+1}{b-1}$

Put this value in (i)

$\left(\frac{b+1}{b-1}\right)^2 + b\left(\frac{b+1}{b-1}\right) - 1 = 0$

$\Rightarrow b^2 + 3b = 0$

$\Rightarrow b(b^2 + 3) = 0$

$\Rightarrow b = 0$ or $b = \pm i\sqrt{3}$

46. Let $\omega \neq 1$ be a cube root of unity and S be the set of all non-singular matrices of the form

$$\begin{bmatrix} 1 & a & b \\ \omega & 1 & c \\ \omega^2 & \omega & 1 \end{bmatrix}$$

where each of a, b and c is either ω or ω^2 . Then the number of distinct matrices in the set S is

- (A) 2 (B) 6 (C) 4 (D) 8

Ans. A

Sol. $\begin{vmatrix} 1 & a & b \\ \omega & 1 & c \\ \omega^2 & \omega & 1 \end{vmatrix} \neq 0$

$(1 - \omega c) - a(\omega - \omega^2 c) + b(\omega^2 - \omega^2) \neq 0$

$1 - \omega c - a\omega + a\omega^2 \neq 0$

$(1 - \omega c) - a\omega(1 - \omega c) \neq 0$

$(1 - \omega c)(1 - a\omega) \neq 0$

$c \neq \omega^2$ & $a \neq \omega^2$ & $b = \omega$ or ω^2

$(a, b, c) \equiv (\omega, \omega, \omega)$ or $(\omega, \omega^2, \omega)$

47. The circle passing through the point $(-1, 0)$ and touching the y-axis at $(0, 2)$ also passes through the point


- (A) $\left(\frac{-3}{2}, 0\right)$ (B) $\left(\frac{-5}{2}, 2\right)$ (C) $\left(\frac{-3}{2}, \frac{5}{2}\right)$ (D) $(-4, 0)$

Ans. D

Sol. $\therefore (h - 0)^2 + (2 - 2)^2 = (h + 1)^2 + (2 - 0)^2$

$h^2 = h^2 + 1 + 2h + 4$

$h = -\frac{5}{2}$



Equation of circle is

$$\left(x + \frac{5}{2}\right)^2 + (y - 2)^2 = \left(-\frac{5}{2} - 0\right)^2$$

$$x^2 + \frac{25}{4} + 5x + y^2 + 4 - 4y = \frac{25}{4}$$

$$x^2 + y^2 + 5x - 4y + 4 = 0$$

from given points only point $(-4, 0)$ satisfies this equation.

48. If $\lim_{x \rightarrow 0} [1 + x \ln(1 + b^2)]^{\frac{1}{x}} = 2b \sin^2 \theta$, $b > 0$ and $\theta \in (-\pi, \pi]$, then the value of θ is

(A) $\pm \frac{\pi}{4}$

(B) $\pm \frac{\pi}{3}$

(C) $\pm \frac{\pi}{6}$

(D) $\pm \frac{\pi}{2}$

Ans. D

Sol. $\lim_{x \rightarrow \infty} [1 + x \ln(1 + b^2)]^{\frac{1}{x}} = 2b \sin^2 \theta$ $b > 0; \theta \in (-\pi, \pi)$

$$\lim_{x \rightarrow \infty} \left([1 + x \ln(1 + b^2)]^{\frac{1}{x \ln(1 + b^2)}} \right)^{\ln(1 + b^2)} = 2b \sin^2 \theta$$

$$e^{\ln(1 + b^2)} = 2b \sin^2 \theta$$

$$1 + b^2 = 2b \sin^2 \theta$$

$$2 \sin^2 \theta = b + \frac{1}{b}$$

$$\text{RHS} = b + \frac{1}{b} \geq 2 \quad \text{as } b > 0$$

$$\text{But LHS} = 2 \sin^2 \theta \leq 2$$

Only possibility

$$2 \sin^2 \theta = 2$$

$$\sin^2 \theta = 1$$

$$\theta = \pm \frac{\pi}{2}$$

Section - II (Total Marks : 16)

(Multiple Correct Answer Type)

This section contains **4 multiple choice questions**. Each question has four choices (A), (B), (C) and (D) out of which **ONE or MORE** is correct.

49. Let E and F be two independent events. The probability that exactly one of them occurs is $\frac{11}{25}$ and the probability of none of them occurring is $\frac{2}{25}$. If P(T) denotes the probability of occurrence of the event T, then

(A) $P(E) = \frac{4}{5}, P(F) = \frac{3}{5}$

(B) $P(E) = \frac{1}{5}, P(F) = \frac{2}{5}$

(C) $P(E) = \frac{2}{5}, P(F) = \frac{1}{5}$

(D) $P(E) = \frac{3}{5}, P(F) = \frac{4}{5}$

Ans. [A] and [D]

Sol. $P(E) (1 - P(F)) + (1 - P(E)) P(F) = \frac{11}{25}$

$P(E) + P(F) - 2P(E) P(F) = \frac{11}{25}$ (1)

$(1 - P(E) - P(F)) P(E) P(F) = \frac{2}{25}$

$1 - P(E) - P(F) + P(E) P(F) = \frac{2}{25}$ (2)

$P(E) + P(E) - P(E) (PF) P(F) = \frac{23}{25}$

From (1) & (2)

$P(E) P(F) = \frac{12}{25}$

And $P(E) + P(F) = \frac{7}{5}$

So either

$P(E) = \frac{4}{5}, P(F) = \frac{3}{5}$ and $P(E) = \frac{3}{5}, P(F) = \frac{4}{5}$

50. If $f(x) = \begin{cases} -x - \frac{\pi}{2}, & x \leq -\frac{\pi}{2} \\ -\cos x, & -\frac{\pi}{2} < x \leq 0 \\ x - 1, & 0 < x \leq 1 \\ \ln x, & x > 1 \end{cases}$

then

(A) $f(x)$ is continuous at $x = -\frac{\pi}{2}$

(B) $f(x)$ is not differentiable at $x = 0$

(C) $f(x)$ is differentiable at $x = 1$

(D) $f(x)$ is differentiable at $x = -\frac{3}{2}$

Ans. [A], [B], [C] and [D]

Sol. At $x = -\frac{\pi}{2}$

LHL = 0, RHL = 0, $f\left(-\frac{\pi}{2}\right) = 0$

So $f(x)$ is continuous at $x = -\frac{\pi}{2}$

At $x = 0$

LHD = 0 ; RHD = 1

So $f(x)$ is not differentiable at $x = 0$

At $x = 1$

LHD = 1, RHD = 1

So $f(x)$ is differentiable at $x = 1$

in $\left(-\frac{\pi}{2}, 0\right)$; $f(x) = -\cos x$

So $f(x)$ is differentiable at $x = -\frac{3}{2}$

51. Let $f : (0, 1) \rightarrow \mathbb{R}$ be defined by $f(x) = \frac{b-x}{1-bx}$

where b is a constant such that $0 < b < 1$. Then

(A) f is not invertible on $(0, 1)$

(B) $f \neq f^{-1}$ on $(0, 1)$ and $f'(b) = \frac{1}{f'(0)}$

(C) $f = f^{-1}$ on $(0, 1)$ and $f'(b) = \frac{1}{f'(0)}$

(D) f^{-1} is differentiable on $(0, 1)$

Ans. [A]

Sol. $f : (0, 1) \rightarrow \mathbb{R}$

$$f(x) = \frac{b-x}{1-bx} \quad \forall b \in (0, 1)$$

$$f'(x) = \frac{b^2-1}{(1-bx)^2} \quad (-)ve$$

So $f(x)$ is monotonically decreasing for $x \in (0, 1)$

So for $x \in (0, 1)$

$$f(x) \in (-1, b)$$

so $f(x)$ is not onto.

So $f(x)$ is not invertible function.

The most appropriate answer to this question is (A, B), but because of ambiguity in language, It has declared (A) as correct answer.

52. Let L be a normal to the parabola $y^2 = 4x$. If L passes through the point $(9, 6)$, then L is given by

(A) $y - x + 3 = 0$

(B) $y + 3x - 33 = 0$

(C) $y + x - 15 = 0$

(D) $y - 2x + 12 = 0$

Ans. [A], [B] and [D]

Sol. $y = mx - 2m - m^3$

It passes through $(9, 6)$

$$6 = 9m - 2m - 2m - m^3$$

$$m^3 - 7m - 6 = 0$$

$$(m - 1)(m - 2)(m + 3) = 0$$

$$\therefore m = -3, 1, 2$$

Hence equation will be

$$y = x - 3, y = 2x - 12 \text{ and } y = -3x + 33$$

Section - III (Total Marks : 24)

(Integer Answer Type)

This section contains **6 questions**. The answer to each of the questions is a **single-digit integer**, ranging 0 to 9. The bubble corresponding to the correct answer is to be darkened in the ORS.

53. The straight line $2x - 3y = 1$ divides the circular region $x^2 + y^2 \leq 6$ into two parts.

If $S = \left\{ \left(2, \frac{3}{4} \right), \left(\frac{5}{2}, \frac{3}{4} \right), \left(\frac{1}{4}, \frac{-1}{4} \right), \left(\frac{1}{8}, \frac{1}{4} \right) \right\}$, then the number of point(s) in S lying inside the smaller part is

Ans. [2]

Sol. Point (x_1, y_1) lies inside the region if $x_1^2 + y_1^2 - 6 \leq 0$ & $2x_1 - 3y_1 - 1 \leq 0$.

$$P_1 \equiv \left(2, \frac{3}{4} \right) \quad 4 + \frac{9}{16} - 6 \geq 0 \quad \text{True}$$

$$4 - \frac{9}{4} - 1 > 0 \quad \text{True}$$

$$P_2 \equiv \left(\frac{5}{2}, \frac{3}{4} \right) \quad \frac{25}{4} + \frac{9}{16} - 6 \leq 0 \quad \text{False}$$


$$P_3 \equiv \left(\frac{1}{4}, \frac{-1}{4} \right) \quad \frac{1}{16} + \frac{1}{16} - 6 \leq 0 \quad \text{True}$$

$$\frac{2}{4} + \frac{3}{4} - 1 > 0 \quad \text{True}$$

$$P_4 \equiv \left(\frac{1}{8}, \frac{1}{4} \right) \quad \frac{1}{64} + \frac{1}{16} - 6 \leq 0 \quad \text{True}$$

$$\frac{2}{8} - \frac{3}{4} - 1 > 0 \quad \text{False}$$

So P_1 & P_3 lies in the interval



54. Let $\omega = e^{\frac{i\pi}{3}}$ and a, b, c, x, y, z be non-zero complex numbers such that

$$a + b + c = x$$

$$a + b\omega + c\omega^2 = y$$

$$a + b\omega^2 + c\omega = z$$

Then the value of $\frac{|x|^2 + |y|^2 + |z|^2}{|a|^2 + |b|^2 + |c|^2}$, is

Ans. [3]

[Note : Question is invalid, but if we take $\omega = e^{\frac{2i\pi}{3}}$ than answer is 3]

Sol. wrong question if $\omega = e^{i2\pi/3}$ then ans is 3. If $w = e^{i\pi/3}$ then no integral solution is possible.

55. The number of distinct real roots of $x^4 - 4x^3 + 12x^2 + x - 1 = 0$ is

Ans. [2]

Sol. Let $f(x) = x^4 - 4x^3 + 12x^2 + x - 1$

Let $\alpha, \beta, \gamma, \delta$ are the root of equation.

$\therefore \alpha\beta\gamma\delta = -1$ so the equation has at least two real roots.(i)

$$f'(x) = 4x^3 - 12x^2 + 24x + 1$$

$$f''(x) = 12x^2 - 24x + 24 = 12((x + 1)^2 + 1)$$

so $f''(x) > 0$ so $f'(x) = 0$ has only one real roots so $f(x) = 0$ has at most two real roots.(ii)

from (i) & (ii)

$f(x) = 0$ has exactly two real roots.

56. Let $y'(x) + y(x)g'(x) = g(x)g'(x)$, $y(0) = 0$, $x \in \mathbb{R}$, where $f'(x)$ denotes $\frac{df(x)}{d(x)}$ and $g(x)$ is a given

non-constant differentiable function on \mathbb{R} with $g(0) = g(2) = 0$. Then the value of $y(2)$ is

Ans. [0]

Sol. $\frac{dy}{dg} + y = g$

$$I. F. = \int 1.dg = g$$

$$y.e^g = \int ge^g .de^g - \int e^g .dg$$

$$ye^g = ge^g - e^g + c$$

$$y = g - 1 + ce^{-g}$$

$$\therefore y(0) = 0 \text{ \& } g(0) = 0$$

at $x = 0$

$$0 = 0 - 1 + Ce^{-0}$$

$$C = 1$$

$$y = g - 1 + e^{-g}$$

at $x = 2$

$$y(2) = 0 - 1 + e^{-0} = 0$$

57. Let M be a 3×3 matrix satisfying $M \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix}$, $M \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$ and $M \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 12 \end{bmatrix}$.

Then the sum of the diagonal entries of M is

Ans. [9]

Sol. Let $M = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$

$\therefore M \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix} \Rightarrow b = -1, e = 2, h = 3$

$\therefore M \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \Rightarrow a = 0, d = 3, g = 2$

$M \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 12 \end{bmatrix} \Rightarrow c = 1, f = -5, i = 7$

So $a + e + i = 0 + 2 + 7 = 9$

58. Let $\vec{a} = -\hat{i} - \hat{k}$, $\vec{b} = -\hat{i} + \hat{j}$ and $\vec{c} = \hat{i} + 2\hat{j} + 3\hat{k}$ be three given vectors. If \vec{r} is a vector such that $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$ and $\vec{r} \cdot \vec{a} = 0$, then the value of $\vec{r} \cdot \vec{b}$, is

Ans. [9]

Sol. $\vec{a} = -\hat{i} - \hat{k}$, $\vec{b} = -\hat{i} + \hat{j}$, $\vec{c} = \hat{i} + 2\hat{j} + 3\hat{k}$

$(\vec{r} - \vec{c}) \times \vec{b} = 0 \Rightarrow \vec{r} - \vec{c} = \lambda \vec{b} \Rightarrow \vec{r} = \vec{c} + \lambda \vec{b}$

$\therefore \vec{r} \cdot \vec{a} = 0$

$\Rightarrow \vec{a} \cdot \vec{c} + \lambda \vec{b} \cdot \vec{a} = 0$

$\Rightarrow \lambda = -\frac{\vec{a} \cdot \vec{c}}{\vec{b} \cdot \vec{a}} = 4$

$\Rightarrow \vec{r} \cdot \vec{b} = \vec{c} \cdot \vec{b} + \lambda |\vec{b}|^2 = 9$

59. Match the statements given in Column I with the intervals/union of intervals given in Column II.

Column-I

Column-II

(A) The set $\left\{ \operatorname{Re} \left(\frac{2iz}{1-z^2} \right) : z \text{ is a complex number, } |z|=1, z \neq \pm 1 \right\}$

(P) $(-\infty, -1) \cup (1, \infty)$

(B) The domain of the function $f(x) = \sin^{-1} \left(\frac{8(3)^{x-2}}{1-3^{2(x-1)}} \right)$, is

(Q) $(-\infty, 0) \cup (0, \infty)$

(C) If $f(\theta) = \begin{vmatrix} 1 & \tan\theta & 1 \\ -\tan\theta & 1 & \tan\theta \\ -1 & -\tan\theta & 1 \end{vmatrix}$,

(R) $[2, \infty)$

then the set $\left\{f(\theta) : 0 \leq \theta < \frac{\pi}{2}\right\}$, is

(S) $(-\infty, -1] \cup [1, \infty)$

(D) If $f(x) = x^3(3x - 10)$, $x \geq 0$ then $f(x)$ is increasing in

(T) $(-\infty, 0] \cup [2, \infty)$

Ans. (A) P, R, S, (B) T, (C) R, (D) R

Sol. (A) Let $z = \cos\theta + i\sin\theta$

so $\frac{2iz}{1-z} = \frac{2i(\cos\theta + i\sin\theta)}{1 - \cos 2\theta - i\sin 2\theta} = -\cos\theta \operatorname{ec}\theta \quad \forall \theta \neq (2n+1)\frac{\pi}{2}$

so $\operatorname{Re}\left(\frac{2iz}{1-z}\right) = -\cos\theta \operatorname{ec}\theta \in (-\infty, -1) \cup [1, \infty)$

(B) $\frac{8 \times 3^{x-2}}{1-3^{2x-2}} = \frac{8 \times 3^x}{9-3^{2x}}$ Let $3^x = t$

So $f(x) = \sin^{-1}\left(\frac{8 \times 3^x}{9-3^{2x}}\right) = \sin^{-1}\left(\frac{8t}{9-t^2}\right)$

$-1 \leq \frac{8t}{9-t^2} \leq 1$ on solving

$x \in (-\infty, 0) \cup [2, \infty) \cup [1]$

(C) $f(\theta) = 2 \sec^2\theta$ so $f(\theta) \in [2, \infty)$

(D) $f(x) = 3x5^{x^2} - 10x3^{x^2}$

$f'(x) = \frac{15\sqrt{x}}{2}(x-2)$

So $f(x)$ is increasing for $f'(x) \leq 0$

$x \in [2, \infty)$

* The most appropriate answer to this question is

$A \rightarrow q ; B \rightarrow p ; C \rightarrow s ; D \rightarrow s$

But because of ambiguity in language, IIT has declared

$A \rightarrow s ; B \rightarrow t ; C \rightarrow r ; D \rightarrow r$ as correct answer.

60. Match the statement given in column-I with the values given in column-II

Column-I

Column-II

(A) If $\vec{a} = \hat{j} + \sqrt{3}\hat{k}$, $\vec{b} = -\hat{j} + \sqrt{3}\hat{k}$ and $\vec{c} = 2\sqrt{3}\hat{k}$ from a triangle,

(P) $\frac{\pi}{6}$

then the internal angle of the triangle between \vec{a} and \vec{b} , is

(Q) $\frac{2\pi}{3}$


(B) If $\int_a^b (f(x) - 3x) dx = a^2 - b^2$, then the value of $f\left(\frac{\pi}{6}\right)$, is

(R) $\frac{\pi}{3}$

(C) The value of $\frac{\pi^2}{\ln 3} \int_{\frac{7}{6}}^{\frac{5}{6}} \sec(\pi x) dx$ is (S) π

(D) The maximum value of $\left| \arg\left(\frac{1}{1-z}\right) \right|$ for $|z| = 1, z \neq 1$ (T) $\frac{\pi}{2}$

Ans. (A) Q, (B) P, (C) S, (D) T]



$$\cos \theta = \frac{-\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = -\frac{1}{2} \Rightarrow \theta = \frac{2\pi}{3}$$

(B) $\int_a^b (f(x) - (x)) dx = a^2 - b^2$

differentiating w.r.t (b).

$$f(b) - 3b = -2b$$

$$f(b) = b$$

So $f\left(\frac{\pi}{6}\right) = \frac{\pi}{6}$; if $a = b$ then any value of $f(x)$ is possible

(C) $I = \frac{\pi^2}{\ln 3} \int_{\frac{7}{6}}^{\frac{5}{6}} \sec \pi x dx$

$$I = \frac{\pi^2}{\ln 3} \left[\ln |\sec \pi x + \tan \pi x| \right]_{\frac{7}{6}}^{\frac{5}{6}}$$

$$I = \frac{\pi}{\ln 3} \cdot \ln 3 = \pi$$

(D) $\therefore |z| = 1$

$$z = \cos \theta + i \sin \theta, \forall \theta \in (-\pi, \pi] \text{ and } \theta \neq 0.$$

$$\left| \operatorname{Arg} \frac{1}{(1-z)} \right| = \left| \operatorname{Arg} \left(\frac{1}{1 - \cos \theta - i \sin \theta} \right) \right| = \left| \operatorname{Arg} \left(\frac{1}{2} + \frac{i \cot \frac{\theta}{2}}{2} \right) \right|$$

$$= \left| \frac{\pi - \theta}{2} \right| \text{ so maximum value is } \pi.$$

* The most appropriate answer to this question is

A \rightarrow q; B \rightarrow p or p, q, r, s & t; C \rightarrow s; D \rightarrow s

But because of ambiguity in language, IIT has declared

A \rightarrow q; B \rightarrow p or p, q, r, s & t; C \rightarrow s; D \rightarrow t as correct answer